

The yeast *Kluyveromyces marxianus* and its biotechnological potential

Gustavo Graciano Fonseca · Elmar Heinzle ·
Christoph Wittmann · Andreas K. Gombert

Received: 28 January 2008 / Revised: 12 March 2008 / Accepted: 13 March 2008 / Published online: 22 April 2008
© Springer-Verlag 2008

Abstract Strains belonging to the yeast species *Kluyveromyces marxianus* have been isolated from a great variety of habitats, which results in a high metabolic diversity and a substantial degree of intraspecific polymorphism. As a consequence, several different biotechnological applications have been investigated with this yeast: production of enzymes (β -galactosidase, β -glucosidase, inulinase, and polygalacturonases, among others), of single-cell protein, of aroma compounds, and of ethanol (including high-temperature and simultaneous saccharification-fermentation processes); reduction of lactose content in food products; production of bioingredients from cheese-whey; bioremediation; as an anticholesterolemic agent; and as a host for heterologous protein production. Compared to its congener and model organism, *Kluyveromyces lactis*, the accumulated knowledge on *K. marxianus* is much smaller and spread over a number of different strains. Although there is no publicly available genome sequence for this species, 20% of the CBS 712 strain genome was randomly sequenced (Llorente et al. in FEBS Lett 487:71–75, 2000). In spite of these facts, *K. marxianus* can envisage a great biotechnological future because of some of its qualities, such as a broad substrate spectrum, thermotolerance, high growth rates, and less tendency to ferment when exposed

to sugar excess, when compared to *K. lactis*. To increase our knowledge on the biology of this species and to enable the potential applications to be converted into industrial practice, a more systematic approach, including the careful choice of (a) reference strain(s) by the scientific community, would certainly be of great value.

Keywords *Kluyveromyces marxianus* ·
Yeast biotechnology · Yeast physiology · Yeast taxonomy

Taxonomic history of the present species *Kluyveromyces marxianus*

Kluyveromyces marxianus was first described in 1888 by E. C. Hansen, which at that time was named *Saccharomyces marxianus* after Marx, the person who originally isolated this yeast from grapes (Lodder and Kreger-van Rij 1952). In their monograph, Lodder and Kreger-van Rij (1952) describe ten strains of *S. marxianus*, among which a particular strain labeled *Zygosaccharomyces marxianus*, which had been deposited at the Centraalbureau voor Schimmelcultures (CBS) in 1922 by H. Schnegg, was arbitrarily chosen as the type strain. This corresponds to the present CBS 712 strain. Some differences among the ten mentioned strains were already pointed out at that time, regarding the formation of pseudomycelium, and the capacities of assimilating and fermenting lactose. Rotting leaves of sisal, sewage of a sugar factory, and “Luftheefe” (aerated yeast) are other habitats from which the strains of *S. marxianus* had been isolated. Already in 1939, Sacchetti had observed that inulin is fermented by *S. marxianus* (Lodder and Kreger-van Rij 1952). Although it was already recognized at that time that *S. marxianus* and *Saccharomyces fragilis*, which had been isolated from kefir in 1909 by

G. G. Fonseca · A. K. Gombert
Department of Chemical Engineering, University of São Paulo,
São Paulo, São Paulo, Brazil

G. G. Fonseca · E. Heinzle · C. Wittmann
Biochemical Engineering Institute, Saarland University,
Saarbrücken, Germany

A. K. Gombert (✉)
Department of Biotechnology, Delft University of Technology,
Julianalaan 67,
2628BC Delft, The Netherlands
e-mail: andreas.gombert@poli.usp.br

Jørgensen, were very closely related, they were considered distinct species (Lodder and Kreger-van Rij 1952). In 1951, Luh and Phaff affirmed that *S. fragilis* “is the only yeast species capable of attacking pectin” (Lodder and Kreger-van Rij 1952). Yoghurt, soft cheese, a lung with tuberculosis, Koumiss (a beverage made of fermented mare’s milk), a human lesion of tonsils and the pharynx, and feces are other habitats from which the 11 strains belonging to the *S. fragilis* species had been obtained from (Lodder and Kreger-van Rij 1952).

Mainly due to differences in spore and ascus morphology, in the capacity of fermenting and oxidizing different sugars, and in the occurrence of hybridization between strains, when compared to true *Saccharomyces* yeasts, there was a need to reclassify the former species *S. fragilis* and *S. marxianus*, besides *Saccharomyces lactis*, into a new taxon (van der Walt 1970). In 1956, van der Walt described the new genus *Kluyveromyces*, the type species of which was *Kluyveromyces polysporus* (van der Walt 1956). Later, it was found that the latter yeast had very similar properties to the three above-mentioned species, and consequently, they were all reclassified into the genus *Kluyveromyces*, which encompassed 18 species in the second edition of *The Yeasts, a taxonomic study* (Lodder 1970). Additional habitats from which strains had been isolated include Bantu Beer, milk of a mastitic cow, asthmatic expecoration, and maize meal (van der Walt 1970). Again, *Kluyveromyces fragilis* was considered closely related but still separate from *K. marxianus*, mainly due to the former’s high capacity of fermenting lactose. Dairy products and human and animal lesions were the prevalent origin of strains in the *K. fragilis* taxon (van der Walt 1970).

In the third edition of *The Yeasts, a taxonomic study* (Kreger-van Rij 1984), the genus *Kluyveromyces* was divided into 11 species. On the basis of interfertility, the taxon *K. marxianus* was organized into seven varieties, which are able to readily hybridize (van der Walt and Johannsen 1984). Concomitantly, the former species *K. fragilis* and *K. lactis* disappeared.

In the most recent edition of *The Yeasts, a taxonomic study* (Kurtzman and Fell 1998), the chapter on the *Kluyveromyces* genus includes 15 species. The seven varieties within the *K. marxianus* species, proposed in the previous edition of the monograph, were eliminated by considering them as either independent species (e.g., *K. lactis* and *K. dobzhanskii*) or synonyms of *K. lactis* or *K. marxianus* (Lachance 1998). This is due to the examination of the genetic structure of populations, in combination with hybridization ability, as criteria for classification. Consequently, the former species or varieties *Kluyveromyces bulgaricus*, *K. cicerisporus*, *K. fragilis*, and *K. wikenii* could not be considered distinct from *K. marxianus* (Lachance 1998).

Since the biological concept of species cannot be applied to homothallic organisms, such as the majority of yeasts in

the *Kluyveromyces* taxon, any classification is always based on arbitrary criteria, which have changed along time, as discussed above. Since the development of rapid and efficient gene sequencing tools, it became natural to utilize gene sequences as the criterion for the comparison and classification of microorganisms into the different taxa. Rather than performing single gene comparisons, the most recent reports on the taxonomy of *Kluyveromyces* yeasts employ multigene sequence analyses for elucidating the phylogeny of the different strains. Using this strategy, Kurtzman and Robnett (2003) showed that the species described by Lachance (1998) in the *Kluyveromyces* genus are actually distributed into six clades, indicating the polyphyly of this group of yeasts. This is mainly due to the previous criteria employed in classification, such as ascus morphology (in this particular case, ascus deliquescence), which are inadequate as phylogenetic descriptors (Kurtzman 2003; Kurtzman and Robnett 2003). It has been proposed that genera should be circumscribed according to the phylogenetically defined clades, rather than on phenotypic analyses (Kurtzman and Robnett 2003). As a result of this, the number of species in the *Kluyveromyces* genus decreased to six and the species *K. marxianus* is proposed as the conserved type species (Kurtzman 2003; Lachance 2007). The type species of the originally described *Kluyveromyces* genus (van der Walt 1956), namely, *K. polysporus*, has been reclassified into the newly proposed *Vanderwaltozyma* genus (Kurtzman 2003; Lachance 2007).

Biochemistry, metabolism, and physiology

It should be noted that the great majority of studies published on *K. marxianus* have not aimed at looking into its biochemistry, metabolism, or physiology. Most of the works that are publicly available explored potential applications of this organism (see “[Biotechnological applications](#)”), without investigating what takes place at the intracellular level. Typically, the yeast cells have been cultivated on a specific substrate, and measurements have been carried out in such a way that only the concentrations of a substrate and of a product, besides the cell concentration, are determined. In what concerns physiology, carbon balances are very rarely looked at, meaning that it is only possible to have a rough macroscopic picture of the cellular reactions and, hence, of the organism’s physiology.

Since the 1970’s, a number of studies has been published on biochemical and metabolic aspects of different *K. marxianus* strains (a summary is presented in Table 1). Some studies were actually aimed at identifying suitable classification methods for *K. marxianus*, which has always been a challenging task. These include the observation that, in contrast to *S. cerevisiae*, ergosterol is the only sterol

Table 1 Compilation of biochemical and metabolic studies performed with the yeast *K. marxianus*

Target of the study	Strain employed	Reference
Sterol composition	<i>K. fragilis</i> NCYC 100	Penman and Duffus (1974)
Subcellular localization of the enzyme alcohol dehydrogenase (ADH)	<i>K. fragilis</i> H32	Künkel and May (1976)
Measurement of heat evolution by microcalorimetry	<i>K. fragilis</i> NCYC 100	Beezer et al. (1979)
Characterization of the enzyme fructose-1,6-bisphosphatase	<i>K. fragilis</i> ATCC 10022	Toyoda and Sy (1984)
Lactose symporter	<i>K. marxianus</i> CBS 397; <i>K. marxianus</i> IGC 2902, CBS 712, IGC 2587, IGC 2671, IGC 3014, NRRL Y-1122, CBS 397	Van den Broek et al. (1987); Carvalho-Silva and Spencer-Martins (1990)
High- and low-affinity glucose transporters	<i>K. marxianus</i> IGC 2587	Gasnier (1987)
Regulation of four transport systems identified during the exponential and the stationary phases of batch growth on glucose	<i>K. marxianus</i> CBS 397	De Bruijne et al. (1988)
High- and low-affinity symporters of glucose and fructose	<i>K. marxianus</i> CBS 6556	Postma and Van den Broek (1990)
Proton-motive force-driven transport of galactose	<i>K. marxianus</i> CBS 397	van Leeuwen et al. (1991)
Transport of lactic acid	<i>K. marxianus</i> IGC 3014	Fonseca et al. (1991)
Mechanism of the enzyme UDP glucose 4-epimerase	<i>K. fragilis</i> ATCC 10022 (presumed, from one of the articles)	Mukherji and Bhaduri (1992); Bhattacharjee and Bhaduri (1992); Ray et al. (1995); Majumdar et al. (1998)
Absence of complex I NADH ubiquinone oxidoreductase	<i>K. marxianus</i> K5 (own collection)	Büschges et al. (1994)
Regulation of adenylate cyclase by Ras proteins	<i>K. marxianus</i> CBS 5795W	Verzotti et al. (1994)
Coenzyme Q system and the monosaccharide pattern of cell wall	<i>K. marxianus</i> (various strains)	Molnár et al. (1996)
Behaviour of <i>K. marxianus</i> during autolysis	<i>K. marxianus</i> CBS 397	Amrane and Prigent (1996)
Presence of killer activity	<i>K. marxianus</i> (isolated as a result of the work)	Abranches et al. (1997)
Glucose repression via Mig1p	<i>K. marxianus</i> SGE11 (Montpellier University)	Cassart et al. (1997)
composition of the cell wall	<i>K. marxianus</i> R157, 1586 (University of New South Wales)	Nguyen et al. (1998)
Presence of active efflux pumps involved in drug resistance	<i>K. marxianus</i> IGC 2671	Prudêncio et al. (2000)
Transport of malic acid via a symport mechanism	<i>K. marxianus</i> ATCC 10022, KMS3 (derivative of CBS 6556)	Queirós et al. (1998)
Identification and characterization of a cell-wall acid phosphatase	<i>K. marxianus</i> Y-610 (identical to ATCC 12424)	Yoda et al. (2000)
Capacity of using xenobiotic compounds as nitrogen source	<i>K. fragilis</i> UU1; <i>K. marxianus</i> IMB3	Ternan and McMullan (2000); Ternan and McMullan (2002)
Response of the NADP ⁺ -dependent glutamate dehydrogenase to nitrogen repression	<i>K. marxianus</i> CBS 6556	de Morais (2003)
Characterization of an amine oxidase	<i>K. marxianus</i> CBS 5795	Corpillo et al. 2003
The transport mechanism of xylose	<i>K. marxianus</i> ATCC 52486	Stambuk et al. 2003

This list does not include enzymes of industrial interest, which are separately listed in Table 2.

present in *K. marxianus*; the use of microcalorimetry for identification purposes; and the characterization of coenzyme Q and monosaccharide patterns of cell walls (Table 1).

In other cases, *K. marxianus* was simply used as the source of specific compounds, which were the actual focus of research (mainly enzymes). Examples of this kind of study include those on fructose-1,6-bisphosphatase, uridine diphosphate (UDP) glucose-4-epimerase, an acid phosphatase, an amine oxidase, protein phosphatases, carboxypeptidases, and aminopeptidases (Tables 1 and 2). A number of transport studies were also carried out, namely, on the transport of sugars (lactose, glucose, fructose, galactose, and xylose) and organic acids (lactic and malic acids) (Table 1). Finally, a few metabolic studies were carried out with the aim of characterizing some non-transport aspects of *K. marxianus*, such as the absence of complex I in the respiratory chain, the regulation of adenylate cyclase by Ras proteins, the functional characterization of Mig1p (involved in glucose repression), analysis of cell-wall composition, presence of efflux pumps and their role in drug resistance, and the regulation of the nicotinamide adenine dinucleotide phosphate (oxidized form) (NADP⁺)-dependent glutamate dehydrogenase (Table 1). In many cases, these studies were performed in parallel with other yeasts, mainly with *S. cerevisiae*.

In terms of biochemical studies on enzymes that have industrial interest, *K. marxianus* has been used as a source of inulinase, β -galactosidase, β -glucosidase, and endopolygalacturonases (Table 2). Besides these, some less widespread enzymes with potential industrial application, such as

protein phosphatases, carboxypeptidases, and aminopeptidases have also been investigated more recently (Table 2).

In the 1970s, physiological studies focusing on the influence of some common environmental factors on the growth of *K. marxianus* started to appear in the literature, as a reflection of the eventual interest in using this yeast for industrial applications. Chassang-Douillet et al. (1973) presented the first clear physiological comparison of *K. marxianus* and *S. cerevisiae*, carried out using synthetic media and demonstrating that the so-called glucose effect was absent in *K. marxianus*, as opposed to *S. cerevisiae*. Later studies reported on the effects of pH (de Sánchez and Castillo 1980), ethanol concentration (Bajpai and Margaritis 1982), and sugar concentration (Margaritis and Bajpai 1983) on the growth kinetics of *K. marxianus*. Importantly, *K. marxianus* started to be included in comparative biochemical and physiological studies on yeast in general, such as those related to catabolite repression (Eraso and Gancedo 1984), sensitivity toward toxins (Sukroongreung et al. 1984), and growth inhibition by fatty acids (Viegas et al. 1989).

The use of defined synthetic media combined with chemostat cultivations for quantitative physiological studies started around the 1990s, with works focusing on the regulation of respiration and fermentation and on the so-called Crabtree-effect in yeasts (van Urk et al. 1990; Verduyn et al. 1992). It was then quantitatively shown that *K. marxianus* presents a strong Crabtree-negative character, since no ethanol production was observed after a glucose pulse applied to respiring cells, in contrast to what is commonly observed with *S.*

Table 2 Studies on the biochemistry of enzymes of industrial interest performed with the yeast *K. marxianus*

Enzyme	Application	Strain	Reference
Inulinase	Production of fructose syrup from inulin-containing feed-stocks	<i>K. fragilis</i> ATCC 12424; <i>K. marxianus</i> CBS 6397, CBS 6556	Workman and Day (1984); Rouwenhorst et al. (1988, 1990a,b)
β -galactosidase	Reduction of lactose content in foods	<i>K. fragilis</i> (several strains); <i>K. marxianus</i> NCYC 111; <i>K. marxianus</i> ATCC 10022; <i>K. marxianus</i> IMB3; <i>K. marxianus</i> CBS 6556	Mahoney et al. (1975); Gonçalves and Castillo (1982); Bacci Júnior et al. (1996); Brady et al. (1995); Martins et al. (2002)
β -glucosidase	Hydrolysis of cellulosic materials	<i>K. fragilis</i> ATCC 12424	Raynal and Guérineau (1984); Leclerc et al. (1987)
Endopolygalacturonases	Reduce of viscosity in fruit processing products	<i>K. marxianus</i> CCT 3172; CCT 3172 (overproducing mutant); an unidentified NCYC isolate	Jia and Wheals (2000)
Protein phosphatases	modification of cheese-making qualities of caseins	<i>K. marxianus</i> (strain not indicated)	Jolivet et al. (2001)
Carboxypeptidases	reduction of bitter taste in protein-containing foods	<i>K. marxianus</i> (own isolate)	Ramírez-Zavala et al. (2004b)
Aminopeptidases	direct processing or aging of dairy and meat products	<i>K. marxianus</i> (own isolate)	Ramírez-Zavala et al. (2004a)

Only studies which focused on the biochemistry of enzymes are indicated here. For studies aiming at enzyme production, please refer to the section on “Biotechnological applications”

cerevisiae and even with *K. lactis*, to a lesser extent (Kiers et al. 1998). This was later confirmed by Bellaver et al. (2004). Castrillo and Ugalde (1993) showed that when oxidoreductive metabolism sets in in *K. marxianus*, as a function of increasing glycolytic flux, the maximum respiratory capacity of the cells has not yet been achieved, which is in contrast with the situation in *S. cerevisiae*, in which the onset of respirofermentative metabolism coincides with the achievement of its maximum respiratory capacity. From these studies, *K. marxianus* was classified as facultatively fermentative and Crabtree-negative (van Dijken et al. 1993). It is important to note that it cannot grow under strictly anaerobic conditions and that the occurrence of ethanol formation is almost exclusively linked to oxygen limitation (Visser et al. 1990; van Dijken et al. 1993; Bellaver et al. 2004). More recently, Blank et al. (2005) showed that *K. marxianus* presents the highest tricarboxylic acid cycle flux during batch growth on glucose among the 14 hemiascomycetous yeasts studied within the Génolevures consortium (Souciet et al. 2000).

Other physiological studies report on various issues, such as flocculation (Fernandes et al. 1992, 1993), the

influence of CO₂ on the survival of *K. marxianus* (Isenschmid et al. 1995), the influence of the specific growth rate on the morphology of the NRRLy2415 strain, which displays significant growth in pseudo-hyphal form (O'Shea and Walsh 2000), the effects of increased air pressure on the biomass yield of *K. marxianus* (Pinheiro et al. 2000), the response of *K. marxianus* to oxidative agents such as hydrogen peroxide (Pinheiro et al. 2002), and the macromolecular composition of *K. marxianus* cells as a function of the specific growth rate (Fonseca et al. 2007). The later data can be particularly useful for metabolic flux analysis studies.

One important aspect on the physiology of *K. marxianus* is the fact that significantly different growth parameters, such as μ_{\max} and $Y_{x/s}$, have been reported not only for different strains within the species but also for the same strain when investigated in different laboratories (Fonseca et al. 2007).

From the data in Tables 1 and 2 (and also in Table 3), it can be observed that the number of strains that have been investigated is quite large, and many of them were not

Table 3 Biotechnologically relevant genes sequenced in *K. marxianus*

Gene/function	Strain	Reference
β -glucosidase	<i>K. fragilis</i> ATCC 12424	Raynal et al. (1987)
<i>INU1</i> /inulinase	<i>K. marxianus</i> ATCC 12424	Laloux et al. (1991)
<i>LEU2</i> / β -isopropylmalate dehydrogenase	<i>K. marxianus</i> CBS 6556	Bergkamp et al. (1991)
<i>URA3</i> /orotidine-5'-phosphate decarboxylase	<i>K. marxianus</i> CBS 6556	Bergkamp et al. (1993b)
<i>PDC</i> /pyruvate decarboxylase	<i>K. marxianus</i> ATCC 10606	Holloway and Sudben (1993)
<i>ADH1</i> /alcohol dehydrogenase	<i>K. marxianus</i> ATCC 12424	Ladrière et al. (1993)
<i>ABF1</i> /a DNA binding protein	<i>K. marxianus</i> (strain not indicated)	Oberyé et al. (1993)
the <i>GAP</i> family/glyceraldehyde-3-phosphate dehydrogenases	<i>K. marxianus</i> ATCC 10022	Fernandes et al. (1995)
<i>LAC4</i> / β -galactosidase	<i>K. fragilis</i> (strains not indicated)	Huo and Li (1995)
<i>MIG1</i> /DNA-binding protein involved in glucose repression	<i>K. marxianus</i> SGE11 (Montpellier University)	Cassart et al. (1997)
<i>EPG1</i> /endopolygalacturonase	<i>K. marxianus</i> BKM Y-719	Šiekštelė et al. (1999)
<i>PCPL3</i> /purine-cytosine permease	<i>K. marxianus</i> ATCC 12424	Ball et al. (1999)
<i>ADH2</i> /alcohol dehydrogenase	<i>K. marxianus</i> ATCC 12424	Ladrière et al. (2000)
17% of the genome (1,300 genes by a partial random strategy)	<i>K. marxianus</i> CBS 712	Llorente et al. (2000)
Inulinase	<i>K. marxianus</i> (strain not indicated)	GenBank AF178979
<i>URA3</i> /orotidine-5'-phosphate decarboxylase	<i>K. cicerisporus</i> CBS 4857	Zhang et al. (2003)
<i>QOR</i> /NADPH quinone oxidoreductase	<i>K. marxianus</i> KCTC 7155	Kim et al. (2003)
<i>URA 9</i> /dihydroorotate dehydrogenase 2	<i>K. marxianus</i> NRRL Y-8281	GenBank AY444339
<i>HIS3</i> /imidazoleglycerol-phosphate dehydratase	<i>K. cicerisporus</i> CBS 4857	GenBank AY303539
<i>OYE</i> /old yellow enzyme	<i>C. macedoniensis</i> AKU 4588	Kataoka et al. (2004)
<i>EPG1-2</i> /endopolygalacturonase	<i>K. marxianus</i> CECT 1043	GenBank AY426825
<i>FPS1</i> /plasma membrane glycerol channel	<i>K. marxianus</i> IGC 3886	Neves et al. (2004)
β -galactosidase	<i>K. marxianus</i> (strain not indicated)	GenBank AY526090
Exoinulinase	<i>K. marxianus</i> IW 9801	GenBank AY649443
<i>TPI1</i> /triosephosphate isomerase	<i>K. marxianus</i> (strain not indicated)	GenBank AJ577476

Partial gene sequences deposited in public databases and sequences coding for RNA were not included. GenBank accession numbers were only given when there is no available article to be cited.

obtained directly from the main culture collections worldwide. If, on the one hand, this leads to an interesting metabolic diversity and to several potential applications, as described below in this review, it makes it difficult, on the other hand, to gain fundamental knowledge on the metabolism and physiology of this yeast. In this sense, it would be necessary that researchers started using a reduced number of strains (chosen from key culture collections), similarly to the way in which the *K. lactis* community has been using the CBS 2359 strain (Lachance 1998; Fukuhara 2006). This would allow the development of efficient molecular genetic tools for *K. marxianus* (probably starting with genome sequencing), which are the basis for performing systematic studies that will finally lead to a better understanding of the biology of this species. A possibility would be to choose one or two strains with characteristics that have given *K. marxianus* a clear advantage over other yeasts: thermotolerance, high growth rate, absence of fermentative metabolism upon sugar excess, and a broad substrate spectrum. For making this choice, an approach as the one reported by van Dijken et al. (2000) could be followed.

Recombinant DNA technology

As with any (potential) industrial organism, rational genetic manipulation is one of the most efficient ways of optimizing process yield and/or productivity. In some cases, the application of recombinant DNA (rDNA) technology may even become a prerequisite for a successful industrial process, either to increase product titer and/or purity to levels at which the process becomes economically feasible or to render the producing host capable of synthesizing a heterologous compound. This kind of activity has been well known as metabolic engineering, which is now a consolidated discipline (Stephanopoulos et al. 1998). rDNA technology is also an invaluable technique for genetic and physiological studies, which in turn are essential for increasing our understanding of *K. marxianus*.

Already more than two decades ago, transformation methods for inserting foreign DNA into *K. marxianus* have been developed. Das et al. (1984) constructed a plasmid called pGL2, containing the kanamycin resistance gene as a dominant selectable marker, and the *KARS2* autonomously replicating sequence of *K. lactis*. They showed that the transformation method of intact cells with alkali cations, originally developed for *Saccharomyces cerevisiae* by Ito et al. (1983), also worked in the strain *K. fragilis* C21. However, the transformation efficiency was rather low.

A breakthrough in molecular biology research of *Kluyveromyces* yeasts was the discovery of the pKD1 plasmid in the species *Kluyveromyces drosophilorum*

(Falcone et al. 1986). The 4.8-kb, 1.65 μ m pKD1 plasmid proved to have a similar organization but different sequences and host specificities, when compared to other already known plasmids, such as the 2 μ plasmid of *Saccharomyces* yeasts (Chen et al. 1986). In contrast to the latter, pKD1 can be maintained stable in *K. lactis*, but not in *S. cerevisiae*, in the absence of selective pressure (Bianchi et al. 1987). Later, it was shown that the insertion of the kanamycin resistance gene, the *URA3* gene of *S. cerevisiae*, a replication origin for *E. coli*, and the ampicillin resistance gene into pKD1 rendered a shuttle plasmid that could be transformed and maintained in *K. marxianus* strains CBS 6556 and CBS 712, though still with low-transformation efficiencies (Chen et al. 1989). This is in accordance with the fact that ARS and centromere sequences of *K. lactis* work in *K. marxianus* and vice versa (Das et al. 1984; Iborra and Ball 1994). Thus, pKD1-based plasmids have become the most common choice for inserting foreign DNA sequences into *K. marxianus* (Bergkamp et al. 1993b; Bartkevičiute et al. 2000; Zhang et al. 2003).

Iborra (1993) reported for the first time transformation efficiencies in the order of hundreds to thousands of transformants per microgram of DNA with *K. marxianus*, either with the lithium method (Ito et al. 1983) or using electroporation (Meilhoc et al. 1990). Similar results were obtained more recently by Zhang et al. (2003).

Besides requiring efficient vectors and transformation protocols, foreign gene expression also depends on the promoter and eventually a signal sequence for directing the synthesized protein into the extracellular environment, which usually facilitates downstream operations. For this purpose, Bergkamp et al. (1993a) used the promoter and prepro-signal sequence of the *INU1* (inulinase) gene to successfully direct heterologous expression and secretion of α -galactosidase in *K. marxianus*, with dramatically higher efficiencies when compared to the use of classical *S. cerevisiae* promoters, such as *PGK*.

With the *INU1* promoter, heterologous gene expression can be fine-tuned by choosing the appropriate carbon source. Another regulated promoter that was successfully used in *K. marxianus* is the tetracycline repressible promoter (Pecota and da Silva 2005).

Strong, constitutive promoters for driving heterologous gene expression have also been described, such as that of a purine-cytosine permease gene (Ball et al. 1999). Instead of fine-tuning foreign gene expression according to promoter strength or induction properties, Pecota et al. (2007) developed an insertion cassette that enables multicopy integration of a precise number of gene copies into *K. marxianus* with recycling of the selection marker.

Auxotrophic mutants of *K. marxianus*, for their use in transformation experiments, have been reported for leucine, uracil, histidine, or tryptophan requirement (Bergkamp et

al. 1991, 1993b; Basabe et al. 1996; Hong et al. 2007). Dominant markers applicable for the selection of *K. marxianus* transformants include at least the kanamycin, the aureobasidin A, and the nurseothricin resistance genes (Das et al. 1984; Hashida-Okado et al. 1998; Goldstein and McCusker 1999; Steensma and Ter Linde 2001; Ribeiro et al. 2007). Recycling of the marker gene for multiple gene disruptions can be performed in *K. marxianus* in the same way as in *S. cerevisiae* and *K. lactis* with the Cre-loxP system (Güldener et al. 1996; Ribeiro et al. 2007).

A number of genes have been cloned and sequenced in different *K. marxianus* strains, and the most relevant ones are indicated in Table 3.

Biotechnological applications

When evaluating the yeast *K. marxianus* for biotechnological applications, it is impossible not to consider other most popular species, mainly *S. cerevisiae* and *K. lactis*. The former is probably the most employed biocatalyst in the biotechnological industry and a model organism in biological studies, whereas the latter has been chosen as a model Crabtree-negative, lactose-utilizing organism (Lachance 1998; Fukuhara 2006). The fact that several *K. marxianus* strains have obtained the generally-regarded-as-safe (GRAS) status, similarly to *S. cerevisiae* and *K. lactis* (Hensing et al. 1995) indicates that this aspect does not impose any disadvantage for the former, when compared to the latter yeasts, in terms of process approval by regulatory agencies. The fact that *K. lactis* was chosen by the scientific community as the model organism in the *Kluyveromyces* genus, and not *K. marxianus*, led not only to a much better understanding of its physiology (for reviews, see e.g., Schaffrath and Breunig 2000; Wolf et al. 2003; Breunig and Steensma 2003 and the whole issue no. 3, vol. 6 of FEMS Yeast Research), and to the full sequencing of its genome (Dujon et al. 2004) but also to the development of several applications, including the expression of more than 40 heterologous proteins (van Ooyen et al. 2006). This is due, to a great extent, to the fact that researchers have used, from the beginning, a very small number of *K. lactis* isolates (Fukuhara 2006), which has not been the case in the *K. marxianus* species.

The development of biotechnological applications with *K. marxianus* has been motivated by a number of advantages it has when compared to *K. lactis*. These include at least the fact that it can grow on a broader variety of substrates and at higher temperatures, its higher specific growth rates, and the lesser tendency to produce ethanol it has when exposed to sugar excess (Rouwenhorst et al. 1988; Steensma et al. 1988; Bellaver et al. 2004; see also “Biochemistry, metabolism, and physiology”).

One very important aspect of the ecology of *K. marxianus* should be taken into account when considering its biotechnological utilization: Individuals have been isolated from an enormous variety of habitats (see “Taxonomic history of the present species *Kluyveromyces marxianus*”).

The obvious consequence is that the metabolic diversity is broad, and hence, potential biotechnological applications of *K. marxianus* strains are manifold. A summary of the most explored applications with this yeast follows.

Although several yeasts have been reported for the production of aroma compounds, only a few of these can find industrial application due to their GRAS status (Medeiros et al. 2000, 2001). *Kluyveromyces* sp. produce aroma compounds such as fruit esters, carboxylic acids, ketones, furans, alcohols, monoterpene alcohols, and isoamyl acetate in liquid fermentation (Scharpf et al. 1986; Fabre et al. 1995). Of all these compounds, 2-phenyl ethanol (2-PE), with rose petals aroma, is the most important commercially (Welsh et al. 1989; Leclercq-Perlat et al. 2004). Natural 2-PE has a high-value (approximately US \$1,000 kg⁻¹) serving a current world market of approximately 7,000 tons per annum (data from 1990; Etschmann et al. 2002). This alcohol presents sensorial characteristics that influence the quality of the wine, distilled drinks, or fermented foods. It is also found in fresh beer and is added to various industrial food products such as ice creams, bullets, non-alcoholic drinks, gelatines, puddings, and bubble gums (Wittmann et al. 2002). The influences of the carbon source (Fabre et al. 1998; Medeiros et al. 2000), aeration rate (Medeiros et al. 2001), media composition (Etschmann et al. 2004), and cultivation conditions (Etschmann and Schrader 2006) on the aroma production using *K. marxianus* were studied.

K. marxianus possesses the natural ability to excrete enzymes. This is a desired property for cost-efficient downstream processing of low- and medium-value enzymes (Hensing et al. 1994). The enzymes that hydrolyze pectic substances are known as pectic enzymes, pectinases, or pectinolytic enzymes (Wimborne and Rickard 1978). Pectinases are industrially used in the extraction and clarification of fruit juices (i.e., grape and apple; Schwan et al. 1997; Blanco et al. 1999). Other interesting applications are related to the maceration of vegetables, oil extraction, and formulation of animal feed using complex mixtures with cellulases to make the nutritional assimilation easier (Blanco et al. 1999). *K. marxianus* has considerable economic advantages over *Aspergillus* as an endo-PG source, even without genetic improvement of the strains (Harsa et al. 1993). Thus, the use of PGs from *K. marxianus* has attracted considerable interest (García-Garibay et al. 1987b; Harsa et al. 1993; Donaghy and McKay 1994). Moreover, no other pectinolytic enzyme, besides PG, was reported to be secreted by *K. marxianus*

CCT 3172 in the culture media, which should facilitate the process to produce pure enzyme (Schwan et al. 1997).

In *K. marxianus*, the pectinolytic enzymes are only produced during exponential growth, but almost all PG is secreted in the start of the stationary phase (Schwan and Rose 1994; Schwan et al. 1997; Serrat et al. 2004). Among the cultivation parameters, dissolved oxygen was reported to be the key in the production of both biomass and endo-PG (Wimborne and Rickard 1978; Garcia-Garibay et al. 1987a). High rates and yields of biomass production require high oxygenation levels that, however, repress endo-PG induction (Wimborne and Rickard 1978). It was reported that *K. marxianus* exhibited pectolytic ability when it was grown without shaking and under anaerobic conditions, and no activity was found at high aeration rates (Barnby et al. 1990; Schwan and Rose 1994; Garcia-Garibay et al. 1987a).

The effect of temperature was assessed on both growth and endo-PG production in combination with the effect of dissolved oxygen (Schwan and Rose 1994). Temperature was also reported to have no direct effect on the synthesis of this enzyme but influenced the growth rate and had an indirect effect due to changes in oxygen solubility (Cruz-Guerrero et al. 1999). Addition of pectin in an aerobic culture in a fermenter was reported to derepress the production of the enzyme (Garcia-Garibay et al. 1987a). However, whereas some authors did not find any effect of pectin addition to the medium in an endo-PG producing strain (Schwan and Rose 1994; Schwan et al. 1997), others reported the enhancement of pectinase production by this yeast when pectin was added (Wimborne and Rickard 1978; Lim et al. 1980; Cruz-Guerrero et al. 1999). *K. marxianus* CCT 3172 was able to break down pectin but required a usable source of carbon and energy to elaborate pectinolytic activity (Schwan and Rose 1994). It had a strong endo-PG activity between pH 4–6 with pH 5 as optimum (Schwan et al. 1997). Furthermore, the type of pectinase excreted by this strain was pointed as a feasible alternative to fungal production due to the lower broth viscosity, which can make downstream operations easier (Almeida et al. 2003a).

Lactose-intolerance can be circumvented by removing lactose from the diet or by converting this sugar into glucose and galactose with β -D-galactosidase (Rajoka et al. 2003). It was reported that only approximately 2% of the recognized yeast species are capable of fermenting lactose (Barnett et al. 1983), among which strains within the *Kluyveromyces* genus can be found. In a screening performed with yeast strains belonging to different genera, only two cultures of *K. fragilis* and *Ferrissia fragilis* showed β -galactosidase activity (Fiedurek and Szczodrak 1994). Lactose is considered the primary inducer of β -D-galactosidase synthesis (Furlan et al. 2000) and production

of lactase by *K. marxianus* using cheese whey as a nutrient source has been investigated by several authors (Sonawat et al. 1981; Nunes et al. 1993).

During the stationary phase of growth, the β -D-galactosidase activity of *K. marxianus* CBS 712 and CBS 6556 remained approximately constant (Rech et al. 1999). In contrast, a reduction of β -D-galactosidase activity was reported by other authors in the stationary phase (Mahoney et al. 1975). The highest β -D-galactosidase activity of *K. marxianus* CBS 712 and CBS 6556 was reported to be at 37°C, decreasing quickly at temperatures above 40°C (Rech et al. 1999), while in *K. marxianus* IMB3, the enzyme is optimally active at 50°C (Barron et al. 1995a). On the other hand, the β -galactosidase of *K. marxianus* CBS 6556 is more stable than the corresponding enzymes of other strains, when stored at low temperatures, e.g., 4°C (Rech et al. 1999; Itoh et al. 1982; Brady et al. 1995). When different substrates were investigated for β -galactosidase production by *K. marxianus*, lactose supported the highest enzyme activities (Rajoka et al. 2003; Rajoka et al. 2004).

Inulinase is an enzyme that cleaves fructose molecules from inulin. Its expression is induced by inulin or sucrose, and the enzyme can be excreted to the culture medium or remain associated to the cell wall (Rouwenhorst et al. 1988; Barranco-Florido et al. 2001). *K. marxianus* has been widely studied for inulinase production, aiming at the production of fructose syrup from inulin (Cruz-Guerrero et al. 1995).

Pessoa and Vitolo (1999) obtained the highest inulinase activities with the *K. marxianus* DSM 70106 strain, using inulin as the carbon source. Growth of *K. marxianus* on sucrose also proceeds via the action of an extracellular inulinase (Hensing et al. 1994), which is repressed when growth is not sucrose-limited (Rouwenhorst et al. 1988; Parekh and Margaritis 1985; Grootwassink and Hewitt 1983).

K. marxianus has also been proposed as a source of: (1) oligonucleotides, used as flavour enhancers in food products; (2) oligosaccharides, used as prebiotics; and (3) oligopeptides, immuno stimulators added to dairy products that are released in the wort after whey protein proteolysis (Belem et al. 1997; Belem and Lee 1998, 1999). Recent studies have shown the potential of *K. marxianus* FII 510700 biomass as an alternative source to *S. cerevisiae* for yeast autolysates (Lukondeh et al. 2003a), alkali-insoluble glucans (Lukondeh et al. 2003c), and a natural bioemulsifier (Lukondeh et al. 2003b).

In the field of bioremediation, processes using *K. marxianus* were developed for the removal of copper ions (II) with molasses as a nutrients source (Aksu and Dönmez 2000). Lead (II) uptake by *K. marxianus* from contaminated molasses had negative effects on cell growth. Nevertheless, the decrease in biomass formation did not lead to decreased lead (II) uptake; on the contrary, the biosorption ability was

higher at higher initial lead (II) concentrations (Skountzou et al. 2003).

Ethanol production at elevated temperatures has received much attention because of the potential cost savings, which could be obtained by continuous evaporation of ethanol from the broth under reduced pressure (Hacking et al. 1984; Gough et al. 1996, 1997, 1998; Banat et al. 1998). This topic was recently reviewed for yeasts in general, including *K. marxianus*. The advantages described, besides the energy savings due to reduced cooling costs, were higher saccharification and fermentation rates, continuous ethanol removal, and reduced contamination (Banat et al. 1998). However, the temperature increase has a negative effect on ethanol yield and also reduces the cell viability (Anderson et al. 1986; Ballesteros et al. 1991). *K. marxianus* was reported to produce alcohol at temperatures above 40°C and to have a maximum growth temperature of 47°C (Anderson et al. 1986), 49°C (Hughes et al. 1984), or even 52°C (Banat et al. 1992). Lower ethanol tolerance was observed when *K. marxianus* was compared to *S. cerevisiae*, and this was correlated with the activity of the plasma membrane ATPase (Rosa and Sa-Correia 1992; Fernanda and Sa-Correia 1992). Hacking et al. (1984) screened yeast strains for their ability to ferment glucose to ethanol at high temperatures. The tolerance of all species seemed to decrease with temperature, but in general, *Kluyveromyces* strains were more thermotolerant than *Saccharomyces*, which in turn can produce higher ethanol yields. Anderson et al. (1986) compared *K. marxianus* strains isolated from sugar mills and CBS strains for ethanol production at high temperatures. The CBS strains produced the same ethanol amounts as the new isolates but with lower cell viability and higher cultivation time. Sakanaka et al. (1996) reported the fusion of a thermotolerant strain of *K. marxianus* with a high ethanol producing strain of *S. cerevisiae*; however, their fermentative capacity was severely impaired and the fusants' thermostability was lower than for either of the parental cells.

While Schwan and Rose (1994) reported that ethanol production in galactose-containing medium was not as high as when glucose was the carbon source, Duvnjak et al. 1987 found that galactose was a better carbon source for ethanol production than glucose; however, the strains employed in both works were different. The conversion of xylose into ethanol by *K. marxianus* was already reported some time ago (Margaritis and Bajpai 1982).

Different process strategies have been used for ethanol production with *K. marxianus*: batch cultures with elevated substrate concentrations (Grubb and Mawson 1993; Barron et al. 1996), fed-batch production (Ferrari et al. 1994; Gough et al. 1998; Love et al. 1996), continuous system (Love et al. 1998), membrane recycle bioreactors (Tin and Mawson 1993), two-stage fermentation (Hack et al. 1994;

Banat et al. 1996), immobilization with β -galactosidase (Hahn-Hägerdal 1985), calcium-alginate-immobilized cells (Bajpai and Margaritis 1987a,b; Marwaha et al. 1988; Nolan et al. 1994; Riordan et al. 1996; Barron et al. 1996; Brady et al. 1996, 1997a,b, 1998; Ferguson et al. 1998; Gough and Mchale 1998), cells immobilized in poly(vinyl alcohol) cryogel beads (Gough et al. 1998), or in Kissiris, a mineral glass foam derived from lava (Nigam et al. 1997; Love et al. 1996, 1998), extractive fed batch cultures (Jones et al. 1993), simultaneous saccharification and fermentation processes with added enzymes (Barron et al. 1995b, 1996, 1997; Boyle et al. 1997; Nilsson et al. 1995; Ballesteros et al. 2002a,b, 2004; Kádár et al. 2004) or by cloning of heterologous cellulase genes (Hong et al. 2007), and the use of mixed cultures (Ward et al. 1995).

Cheese whey contains lactose and a protein fraction sufficiently rich in essential amino acids. Cheese whey cultivations with *K. marxianus* have been proposed, with promising results, as a means of reducing the pollution caused by this industrial waste stream (Ghaly and Singh 1989; Giec and Kosikowski 1992; Harden 1996; Aktas et al. 2005) and/or to produce single-cell protein (Giec and Kosikowski 1992; Ben-Hassan et al. 1992, Ben-Hassan and Ghaly 1995; Belem and Lee 1999; Schultz et al. 2006; Ghaly and Kamal 2004). Aerobic cultures of microorganisms in cheese whey can reduce up to 90–95% of its BOD (Grubb and Mawson 1993), resulting in bioingredients of high added value for the food industry (Belem et al. 1997).

Other potential applications of the yeast *K. marxianus*, which can be found in the literature include its use as baker's yeast (Caballero et al. 1995) and as an anticholesterolemic agent (Yoshida et al. 2004). The cellular components involved in the hypocholesterolemic activity of *K. marxianus* were further examined (Yoshida et al. 2005).

Last but not the least, *K. marxianus* has been investigated as a host for the production of heterologous proteins. In general, yeasts are capable of performing some post-translational modifications of proteins, such as glycosylation and/or other modifications required for optimal biological activity and stability (Hensing et al. 1995). *S. cerevisiae* has been the most commonly used yeast host for the production of heterologous proteins (Romanos et al. 1992; Gellissen and Hollenberg 1997; Porro et al. 2005). Nevertheless, this yeast has some drawbacks, such as its strong aerobic fermentation behavior and a tendency to hyperglycosylate secreted glycoproteins (Hensing et al. 1994). *K. lactis* has also been used for the production of heterologous proteins (van den Berg et al. 1990; Panuwatsuk and da Silva 2002; Bartkevičiute and Sasnauskas 2003; van Ooyen et al. 2006). *K. marxianus*, which is phylogenetically close to *K. lactis*, is supposed to have a similar capacity for synthesis and secretion of high molecular weight proteins (Wésolowski-Louvel et al. 1996). Some

examples showing that heterologous protein production in this yeast is possible have been reported in the literature (Bergkamp et al. 1993a; Bartkevičiūtė et al. 2000; Zhang et al. 2003). More recently, Pecota et al. (2007) successfully expressed lactate dehydrogenase activity in *K. marxianus*, using an integrative multi-copy system, resulting in lactate production by this yeast. Hong et al. (2007) expressed thermostable endo- β -1,4-glucanase, cellobiohydrolase, and β -glucosidase, also making use of an integrative system, generating a strain capable of converting cellulosic materials into ethanol. Although these studies demonstrate that the heterologous proteins expressed in *K. marxianus* were functional, the capacity of *K. marxianus* to perform post-translational modifications of heterologous proteins still remains to be investigated.

Acknowledgements Grants from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) (Brazil), Deutscher Akademischer Austausch Dienst (DAAD) (Germany), and Fundação Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) (Brazil) are acknowledged.

References

- Abranches J, Morais PB, Rosa CA, Mendonça-Hagler LC, Hagler AN (1997) The incidence of killer activity and extracellular proteases in tropical yeast communities. *Can J Microbiol* 43:328–336
- Aksu Z, Dönmez G (2000) The use of molasses in copper (II) containing wastewaters: effects on growth and copper (II) bioaccumulation properties of *Kluyveromyces marxianus*. *Proc Biochem* 36:451–458
- Aktas N, Boyacı IH, Mutlu M, Tanyolac A (2005) Optimization of lactose utilization in deproteinated whey by *Kluyveromyces marxianus* using response surface methodology (RSM). *Biores Technol* 97:2252–2259
- Almeida C, Branyik T, Moradas-Ferreira P, Teixeira J (2003a) Continuous production of pectinase by immobilized yeast cells on spent grains. *J Biosci Bioeng* 96:513–518
- Amrane A, Prigent Y (1996) Behaviour of the yeast *Kluyveromyces marxianus* var. *marxianus* during its autolysis. *Antonie van Leeuwenhoek* 69:267–272
- Anderson PJ, McNeil K, Watson K (1986) High-efficiency carbohydrate fermentation to ethanol at temperatures above 40°C by *Kluyveromyces marxianus* var. *marxianus* isolated from sugar mills. *Appl Environ Microbiol* 51:1314–1320
- Bacci Júnior M, Siqueira CG, Antoniazi SA, Ueta J (1996) Location of the β -galactosidase of the yeast *Kluyveromyces marxianus* var. *marxianus* ATCC 10022. *Antonie van Leeuwenhoek* 69:357–361
- Bajpai P, Margaritis A (1982) Ethanol inhibition kinetics of *Kluyveromyces marxianus* grown on Jerusalem artichoke juice. *Appl Environ Microbiol* 44:1325–1329
- Bajpai P, Margaritis A (1987a) Kinetics of ethanol production by immobilized *Kluyveromyces marxianus* cells at varying sugar concentrations of Jerusalem artichoke juice. *Appl Microbiol Biotechnol* 26:447–449
- Bajpai P, Margaritis A (1987b) The effect of temperature and pH on ethanol production by free and immobilized cells of *Kluyveromyces marxianus* grown on Jerusalem artichoke extract. *Bio-technol Bioeng* 30:306–313
- Ball MM, Raynal A, Guerineau M, Iborra F (1999) Construction of efficient centromeric, multicopy and expression vectors for the yeast *Kluyveromyces marxianus* using homologous elements and the promoter of a purine-cytosine-like permease. *J Mol Microbiol Biotechnol* 1:347–353
- Ballesteros I, Ballesteros M, Cabanas A, Carrasco J, Martín C, Negro MJ, Saez F, Saez R (1991) Selection of thermotolerant yeasts for simultaneous saccharification and fermentation (SSF) of cellulose to ethanol. *Appl Biochem Biotechnol* 28:307–315
- Ballesteros I, Oliva JM, Negro MJ, Manzanares P, Ballesteros M (2002a) Simultaneous saccharification and fermentation process for converting the cellulosic fraction of olive oil extraction residue into ethanol. *Grasas y Aceites* 53:282–288
- Ballesteros M, Oliva JM, Manzanares P, Negro MJ, Ballesteros I (2002b) Ethanol production from paper material using a simultaneous saccharification and fermentation system in a fed-batch basis. *World J Microbiol Biotechnol* 18:559–561
- Ballesteros M, Oliva JM, Negro MJ, Manzanares P, Ballesteros I (2004) Ethanol from lignocellulosic materials by a simultaneous saccharification and fermentation process (SFS) with *Kluyveromyces marxianus* CECT 10875. *Proc Biochem* 39:1843–1848
- Banat IM, Nigam P, Marchant R (1992) Isolation of thermotolerant, fermentative yeasts growing at 52°C and producing ethanol at 45°C and 50°C. *World J Microbiol Biotechnol* 8:259–263
- Banat IM, Singh D, Marchant R (1996) The use of a thermotolerant fermentative *Kluyveromyces marxianus* IMB3 yeast strain for ethanol production. *Acta Biotechnol* 16:215–223
- Banat IM, Nigam P, Singh D, Marchant R, McHale AP (1998) Ethanol production at elevated temperatures and alcohol concentrations: Part I - Yeasts in general. *World J Microbiol Biotechnol* 14:809–821
- Barnby FM, Morpeth FF, Pyle DL (1990) Endopolygalacturonase production from *Kluyveromyces marxianus*. Resolution, purification, and partial characterisation of the enzyme. *Enzyme Microb Technol* 12:891–897
- Barnett JA, Payne RW, Yarrow D (1983) Yeasts: characteristics and identification. Cambridge University Press, Cambridge
- Barranco-Florido E, García-Garibay M, Gómez-Ruiz L, Azaola A (2001) Immobilization system of *Kluyveromyces marxianus* cells in barium alginate for inulin hydrolysis. *Proc Biochem* 37:513–519
- Barron N, Marchant R, McHale L, McHale AP (1995a) Partial characterization of β -glucosidase activity produced by *Kluyveromyces marxianus* IMB3 during growth on cellobiose containing media at 45°C. *Biotechnol Lett* 17:1047–1050
- Barron N, Marchant R, McHale L, McHale AP (1995b) Studies on the use of a thermotolerant strain of *Kluyveromyces marxianus* in simultaneous saccharification and ethanol formation from cellulose. *Appl Microbiol Biotechnol* 43:518–520
- Barron N, Marchant R, McHale L, McHale AP (1996) Ethanol production from cellulose at 45°C using a batch-fed system containing alginate-immobilized *Kluyveromyces marxianus* IMB3. *World J Microbiol Biotechnol* 12:103–104
- Barron N, Mulholland H, Boyle M, McHale AP (1997) Ethanol production by *Kluyveromyces marxianus* IMB3 during growth on straw-supplemented whiskey distillery spentwash at 45°C. *Bioproc Eng* 17:383–386
- Bartkevičiūtė D, Sasnauskas K (2003) Studies of yeast *Kluyveromyces lactis* mutations conferring super-secretion of recombinant proteins. *Yeast* 20:1–11
- Bartkevičiūtė D, Šiekštelė R, Sasnauskas K (2000) Heterologous expression of the *Kluyveromyces marxianus* endopolygalacturonase gene (*EPG1*) using versatile autonomously replicating vector for a wide range of host. *Enzyme Microb Technol* 26:653–656
- Basabe L, Cabrera N, Yong V, Menéndez J, Delgado JM, Rodríguez L (1996) Isolation and characterization of mutants as an approach to a transformation system in *Kluyveromyces marxianus*. *Curr Genet* 30:89–92

- Beezer AE, Newell RD, Tyrrell HJ (1979) Characterisation and metabolic studies of *Saccharomyces cerevisiae* and *Kluyveromyces fragilis* by flow microcalorimetry. *Antonie van Leeuwenhoek* 45:55–63
- Belem MAF, Lee BH (1998) Production of bioingredients from *Kluyveromyces marxianus* grown on whey: an alternative. *Crit Rev Food Sci Nut* 38:565–598
- Belem MAF, Lee BH (1999) Fed-batch fermentation to produce oligonucleotides from *Kluyveromyces marxianus* grown on whey. *Proc Biochem* 34:501–509
- Belem MAF, Gibbs BF, Lee BH (1997) Enzymatic production of ribonucleotides from autolysates of *Kluyveromyces marxianus* grown on whey. *J Food Sci* 62:851–857
- Bellaver LH, de Carvalho NMB, Abrahão-Neto J, Gombert AK (2004) Ethanol formation and enzyme activities around glucose-6-phosphate in *Kluyveromyces marxianus* CBS 6556 exposed to glucose or lactose excess. *FEMS Yeast Res* 4:691–698
- Ben-Hassan RM, Ghaly AE (1995) Continuous production of single-cell protein from cheese whey lactose using *Kluyveromyces fragilis*. *Trans ASAE* 38:1121–1127
- Ben-Hassan RM, Ghaly AE, Ben-Abdallah N (1992) Metabolism of cheese whey lactose by *Kluyveromyces fragilis* for energy and growth under batch condition. *Appl Biochem Biotechnol* 33:97–116
- Bergkamp RJ, Geerse RH, Verbakel JM, Musters W, Planta RJ (1991) Cloning and disruption of the *LEU2* gene of *Kluyveromyces marxianus* CBS 6556. *Yeast* 7:963–970
- Bergkamp RJM, Bootsma TC, Toschka HY, Mooren ATA, Kox L, Verbakel JMA, Geerse RH, Planta RJ (1993a) Expression of an α -galactosidase gene under control of the homologous inulinase promoter in *Kluyveromyces marxianus*. *Appl Microbiol Biotechnol* 40:309–317
- Bergkamp RJ, Geerse RH, Verbakel JM, Planta RJ (1993b) Cloning and sequencing of the *URA3* gene of *Kluyveromyces marxianus* CBS 6556. *Yeast* 9:677–681
- Bhattacharjee H, Bhaduri A (1992) Distinct functional roles of two active site thiols in UDPglucose 4-epimerase from *Kluyveromyces fragilis*. *J Biol Chem* 267:11714–11720
- Bianchi MM, Falcone C, Re CX, Węslowski-Louvel M, Frontali L, Fukuhara H (1987) Transformation of the yeast *Kluyveromyces lactis* by new vectors derived from the 1.6 μ m circular plasmid pKD1. *Curr Genet* 12:185–192
- Blanco P, Sieiro C, Villa TG (1999) Production of pectic enzymes in yeasts. *FEMS Microbiol Lett* 175:1–9
- Blank LM, Lehmbek F, Sauer U (2005) Metabolic-flux and network analysis in fourteen hemiascomycetous yeasts. *FEMS Yeast Res* 5:545–558
- Boyle M, Barron N, McHale AP (1997) Simultaneous saccharification and fermentation of straw to ethanol using the thermotolerant yeast strain *Kluyveromyces marxianus* IMB 3. *Biotechnol Lett* 19:49–51
- Brady D, Marchant R, McHale L, McHale AP (1995) Isolation and partial characterization of β -galactosidase activity produced by a thermotolerant strain of *Kluyveromyces marxianus* during growth on lactose-containing media. *Enzyme Microb Technol* 17:696–699
- Brady D, Nigam P, Marchant R, McHale L, McHale AP (1996) Ethanol production at 45°C by *Kluyveromyces marxianus* IMB3 immobilized in magnetically responsive alginate matrices. *Biotechnol Lett* 18:1213–1216
- Brady D, Nigam P, Marchant R, McHale AP (1997a) Ethanol production at 45°C by alginate-immobilized *Kluyveromyces marxianus* IMB3 during growth on lactose-containing media. *Bioproc Eng* 16:101–104
- Brady D, Nigam P, Marchant R, Singh D, McHale AP (1997b) The effect of Mn^{2+} on ethanol production from lactose using *Kluyveromyces marxianus* IMB3 immobilized in magnetically responsive matrices. *Bioproc Eng* 17:31–34
- Brady D, Logan SR, McHale AP (1998) The effect of soluble alginate and calcium on β -galactosidase activity produced by the thermotolerant, ethanol-producing yeast strain *Kluyveromyces marxianus* IMB3. *Bioproc Eng* 18:101–104
- Breunig KD, Steensma HY (2003) *Kluyveromyces lactis*: genetics, physiology, and application. In: de Winde JH (ed) *Functional genetics of industrial yeasts*, Topics in current genetics, vol. 2. Springer-Verlag, Berlin Heidelberg New York, pp 171–205
- Büschges R, Bahrenberg G, Zimmermann M, Wolf K (1994) NADH: ubiquinone oxidoreductase in obligate aerobic yeasts. *Yeast* 10:475–479
- Caballero R, Olguín P, Cruz-Guerrero A, Gallardo F, García-Garibay M, Gómez-Ruiz L (1995) Evaluation of *Kluyveromyces marxianus* as baker's yeast. *Food Res Int* 28:37–41
- Carvalho-Silva M, Spencer-Martins I (1990) Modes of lactose uptake in the yeast species *Kluyveromyces marxianus*. *Antonie van Leeuwenhoek* 57:77–81
- Cassart JP, Ostling J, Ronne H, Vandenhoute J (1997) Comparative analysis in three fungi reveals structurally and functionally conserved regions in the Mig1 repressor. *Mol Gen Genet* 255:9–18
- Castrillo JI, Ugalde UO (1993) Patterns of energy metabolism and growth kinetics of *Kluyveromyces marxianus* in whey chemostat culture. *Appl Microbiol Biotechnol* 40:386–393
- Chassang-Douillet A, Ladet J, Boze H, Galzy P (1973) Respiratory metabolism of *Kluyveromyces fragilis* van der Walt. *Z Allg Mikrobiol* 13:193–199
- Chen XJ, Saliola M, Falcone C, Bianchi MM, Fukuhara H (1986) Sequence organization of the circular plasmid pKD1 from the yeast *Kluyveromyces drosophilum*. *Nucleic Acids Res* 14:4471–4481
- Chen XJ, Bianchi MM, Suda K, Fukuhara H (1989) The host range of the pKD1-derived plasmids in yeast. *Curr Genet* 16:95–98
- Corpillo D, Valetti F, Giuffrida MG, Conti A, Rossi A, Finazzi-Agrò A, Giunta C (2003) Induction and characterization of a novel amine oxidase from the yeast *Kluyveromyces marxianus*. *Yeast* 20:369–379
- Cruz-Guerrero A, García-Peña I, Bárzana E, García-Garibay M, Gómez-Ruiz L (1995) *Kluyveromyces marxianus* CDBB-L-278: a wild inulinase hyperproducing strain. *J Ferm Bioeng* 80:159–163
- Cruz-Guerrero A, Bárzana E, García-Garibay M, Gómez-Ruiz L (1999) Dissolved oxygen threshold for the repression of endopolygalacturonase production by *Kluyveromyces marxianus*. *Proc Biochem* 34:621–624
- Das S, Kellermann E, Hollenberg CP (1984) Transformation of *Kluyveromyces fragilis*. *J Bacteriol* 158:1165–1167
- de Bruijne AW, Schuddemat J, van den Broek PJA, van Steveninck J (1988) Regulation of sugar transport systems of *Kluyveromyces marxianus*: the role of carbohydrates and their catabolism. *Biochim Biophys Acta* 939:569–576
- de Moraes MA Jr (2003) The NADP⁺-dependent glutamate dehydrogenase of the yeast *Kluyveromyces marxianus* responds to nitrogen repression similarly to *Saccharomyces cerevisiae*. *Braz J Microbiol* 34:334–338
- de Sánchez SB, Castillo FJ (1980) Effect of pH on the growth of *Kluyveromyces fragilis* on deproteinized whey. *Acta Cient Venez* 31:24–26
- Donaghy JA, McKay AM (1994) The use of *Kluyveromyces fragilis* for the extraction of orange peel pectins. *J Appl Bacteriol* 76:506–510
- Dujon B, Sherman D, Fischer G et al (2004) Genome evolution in yeasts. *Nature* 430:35–44
- Duvnjak Z, Houle C, Mok KL (1987) Production of ethanol and biomass from various carbohydrates by *Kluyveromyces fragilis*. *Biotechnol Lett* 9:343–346

- Eraso P, Gancedo JM (1984) Catabolite repression in yeasts is not associated with low levels of cAMP. *Eur J Biochem* 141:195–198
- Etschmann MMW, Schrader J (2006) An aqueous-organic two-phase bioprocess for efficient production of the natural aroma chemicals 2-phenylethanol and 2-phenylethylacetate with yeast. *Appl Microbiol Biotechnol* 71:440–443
- Etschmann MMW, Bluemke W, Sell D, Schrader J (2002) Biotechnological production of 2-phenylethanol. *Appl Microbiol Biotechnol* 59:1–8
- Etschmann MMW, Sell D, Schrader J (2004) Medium optimization for the production of the aroma compound 2-phenylethanol using a genetic algorithm. *J Mol Catal B: Enzym* 29:187–193
- Fabre CE, Duviau VJ, Blanc PJ, Goma G (1995) Identification of volatile flavour compounds obtained in culture of *Kluyveromyces marxianus*. *Biotechnol Lett* 17:1207–1212
- Fabre CE, Blanc PJ, Goma G (1998) Production of 2-phenylethyl alcohol by *Kluyveromyces marxianus*. *Biotechnol Prog* 14:270–274
- Falcone C, Saliola M, Chen XJ, Frontali L, Fukuhara H (1986) Analysis of a 1.6-micron circular plasmid from the yeast *Kluyveromyces drosophilorum*: structure and molecular dimorphism. *Plasmid* 15:248–252
- Ferguson P, Mulholland H, Barron N, Brady D, McHale AP (1998) Sucrose-supplemented distillery spent-wash as a medium for production of ethanol at 45°C by free and alginate-immobilized preparations of *Kluyveromyces marxianus* IMB3. *Bioproc Eng* 18:257–259
- Fernanda R, Sa-Correia I (1992) Ethanol tolerance and activity of plasma membrane ATPase in *Kluyveromyces marxianus* and *Saccharomyces cerevisiae*. *Enzyme Microb Technol* 14:23–27
- Fernandes PA, Keen JN, Findlay JBC, Moradas-Ferreira PA (1992) Protein homologous to glyceraldehyde-3-phosphate dehydrogenase is induced in the cell wall of a flocculant *Kluyveromyces marxianus*. *Biochim Biophys Acta* 1159:67–73
- Fernandes PA, Sousa M, Moradas-Ferreira P (1993) Flocculation of *Kluyveromyces marxianus* is induced by a temperature upshift. *Yeast* 9:859–866
- Fernandes PA, Sena-Esteves M, Moradas-Ferreira P (1995) Characterization of the glyceraldehyde-3-phosphate dehydrogenase gene family from *Kluyveromyces marxianus* - polymerase chain reaction single-strand conformation polymorphism as a tool for the study of multigenic families. *Yeast* 11:725–733
- Ferrari MD, Loperena L, Varela H (1994) Ethanol production from concentrated whey permeate using a fed-batch culture of *Kluyveromyces fragilis*. *Biotechnol Lett* 16:205–210
- Fiedurek J, Szczodrak J (1994) Selection of strain, culture conditions and extraction procedures for optimum production of b-galactosidase from *Kluyveromyces fragilis*. *Acta Microbiol Pol* 43:57–65
- Fonseca A, Spencer-Martins I, van Uden N (1991) Transport of lactic acid in *Kluyveromyces marxianus*: evidence for a monocarboxylate uniport. *Yeast* 7:775–780
- Fonseca GG, Gombert AK, Heinze E, Wittmann C (2007) Physiology of the yeast *Kluyveromyces marxianus* during batch and chemostat cultures with glucose as the sole carbon source. *FEMS Yeast Res* 7:422–435
- Fukuhara H (2006) *Kluyveromyces lactis* - A retrospective. *FEMS Yeast Res* 6:323–324
- Furlan SA, Schneider ALS, Merkle R, Carvalho-Jonas MD, Jonas R (2000) Formulation of a lactose-free, low-cost culture medium for the production of b-D-galactosidase by *Kluyveromyces marxianus*. *Biotechnol Lett* 22:589–593
- Garcia-Garibay M, Gómez-Ruiz L, Bázquez E (1987a) Studies on the simultaneous production of single-cell protein and endopolygalacturonase from *Kluyveromyces fragilis*. *Biotechnol Lett* 9:411–416
- Garcia-Garibay M, Torres J, López-Munguía-Canales A, Casas LT (1987b) Influence of oxygen transfer rate on b-galactosidase production from *Kluyveromyces marxianus*. *Biotechnol Lett* 9:417–420
- Gasnier B (1987) Characterization of low- and high-affinity glucose transports in the yeast *Kluyveromyces marxianus*. *Biochim Biophys Acta* 903:425–433
- Gellissen G, Hollenberg CP (1997) Application of yeasts in gene expression studies: A comparison of *Saccharomyces cerevisiae*, *Hansenula polymorpha* and *Kluyveromyces lactis*—a review. *Gene* 190:87–97
- Ghaly AE, Singh RK (1989) Pollution potential reduction of cheese whey through yeast fermentation. *Appl Biochem Biotechnol* 22:181–203
- Ghaly AE, Kamal M (2004) Submerged yeast fermentation of acid cheese whey for protein production and pollution potential reduction. *Water Res* 38:631–644
- Giec A, Kosikowski FV (1992) Activity of lactose fermenting yeasts in producing biomass from concentrated whey permeates. *J Food Sci* 47:1892–1894
- Gonçalves JA, Castillo FJ (1982) Partial Purification and characterization of b-D-galactosidase from *Kluyveromyces marxianus*. *J Dairy Sci* 65:2088–2094
- Gough S, McHale AP (1998) Continuous ethanol production from molasses at 45°C using alginate-immobilized *Kluyveromyces marxianus* IMB3 in a continuous-flow bioreactor. *Bioproc Eng* 19:33–36
- Gough S, Flynn O, Hack CJ, Marchant R (1996) Fermentation of molasses using a thermotolerant yeast, *Kluyveromyces marxianus* IMB3: simplex optimization of media supplements. *Appl Microbiol Biotechnol* 46:187–190
- Gough S, Brady D, Nigam P, Marchant R, McHale AP (1997) Production of ethanol from molasses at 45°C using alginate-immobilized *Kluyveromyces marxianus* IMB3. *Bioproc Eng* 16:389–392
- Gough S, Barron N, Zubov AL, Lozinsky VI, McHale AP (1998) Production of ethanol from molasses at 45°C using *Kluyveromyces marxianus* IMB3 immobilized in calcium alginate gels and poly(vinyl alcohol) cryogel. *Bioproc Eng* 19:87–90
- Grootwassink JWD, Hewitt GM (1983) Inducible and constitutive formation of b-fructofuranosidase (inulase) in batch and continuous cultures of the yeast *Kluyveromyces marxianus*. *J Gen Microbiol* 129:31–41
- Grubb CF, Mawson AJ (1993) Effects of elevated solute concentrations on the fermentation of lactose by *Kluyveromyces marxianus* Y-113. *Biotechnol Lett* 15:621–626
- Goldstein AL, McCusker JH (1999) Three new dominant drug resistance cassettes for gene disruption in *Saccharomyces cerevisiae*. *Yeast* 15:1541–1553
- Güldener U, Heck S, Fielder T, Beinhauer J, Hegemann JH (1996) A new efficient gene disruption cassette for repeated use in budding yeast. *Nucleic Acids Res* 24:2519–2524
- Hack CJ, Banat IM, Singh D, Marchant R (1994) Ethanol production by a strain of *Kluyveromyces marxianus* at elevated temperatures in various bioreactor configurations. In: Proceedings of Conference on Fermentation Physiology, Pub. Institution of Chemical Engineers, Brighton, pp 7–9
- Hacking AJ, Taylor IWF, Hanas CM (1984) Selection of yeasts able to produce ethanol from glucose at 40°C. *Appl Microbiol Biotechnol* 19:361–363
- Hahn-Hägerdal B (1985) Comparison between immobilized *Kluyveromyces fragilis* and *Saccharomyces cerevisiae* coimmobilized with b-galactosidase, with respect to continuous ethanol production from concentrated whey permeate. *Biotechnol Bioeng* 27:914–916
- Harden TJ (1996) The reduction of BOD and production of biomass from acid whey by *Kluyveromyces marxianus*. *Food Aust* 48:456–457

- Harsa S, Zaror CA, Pyle DL (1993) Production of polygalacturonases from *Kluyveromyces marxianus* fermentation—preliminary process design and economics. *Proc Biochem* 28:187–195
- Hashida-Okado T, Ogawa A, Kato I, Takesako K (1998) Transformation system for prototrophic industrial yeasts using the *AURI* gene as a dominant selection marker. *FEBS Lett* 425:117–122
- Hensing M, Vrouwenvelder H, Hellinga C, Baartmans R, van Dijken JP (1994) Production of extracellular inulinase in high-cell-density fed-batch cultures of *Kluyveromyces marxianus*. *Appl Microbiol Biotechnol* 42:516–521
- Hensing MC, Rouwenhorst RJ, Heijnen JJ, van Dijken JP, Pronk JT (1995) Physiological and technological aspects of large-scale heterologous-protein production with yeasts. *Antonie van Leeuwenhoek* 67:261–279
- Holloway P, Subden RE (1993) The isolation and nucleotide sequence of the pyruvate decarboxylase gene from *Kluyveromyces marxianus*. *Curr Genet* 24:274–277
- Hong J, Wang Y, Kumagai H, Tamaki H (2007) Construction of thermotolerant yeast expressing thermostable cellulase genes. *J Biotechnol* 130:114–123
- Hughes DB, Tudrosen NJ, Moye CJ (1984) The effect of temperature on the kinetics of ethanol production by a thermotolerant strain of *Kluyveromyces marxianus*. *Biotechnol Lett* 6:1–6
- Huo K, Li Y (1995) Cloning and expression of *Kluyveromyces fragilis LAC4* gene. *Sci China B* 38:1332–1340
- Iborra F (1993) High efficiency transformation of *Kluyveromyces marxianus* by a replicative plasmid. *Curr Genet* 24:181–183
- Iborra F, Ball MM (1994) *Kluyveromyces marxianus* small DNA fragments contain both autonomous replicative and centromeric elements that also function in *Kluyveromyces lactis*. *Yeast* 10:1621–1629
- Isenschmid A, Marison IW, von Stockar U (1995) The influence of pressure and temperature of compressed CO₂ on the survival of yeast cells. *J Biotechnol* 39:229–237
- Ito H, Fukuda Y, Murata K, Kimura A (1983) Transformation of intact yeast cells treated with alkali cations. *J Bacteriol* 153:163–168
- Itoh T, Susuki M, Adachi S (1982) Production of β -galactosidase from lactose-fermenting yeasts. *Agric Biol Chem* 46:899–904
- Jia J, Wheals A (2000) Endopolygalacturonase genes and enzymes from *Saccharomyces cerevisiae* and *Kluyveromyces marxianus*. *Curr Genet* 38:264–270
- Jolivet P, Bergeron E, Benyair H, Meunier JC (2001) Characterization of major protein phosphatases from selected species of *Kluyveromyces*. Comparison with protein phosphatases from *Yarrowia lipolytica*. *Can J Microbiol* 47:861–870
- Jones TD, Havard JM, Daugulis AJ (1993) Ethanol production from lactose by extractive fermentation. *Biotechnol Lett* 15:871–876
- Kádár ZS, Szengyel ZS, Réczey K (2004) Simultaneous saccharification and fermentation (SSF) of industrial wastes for the production of ethanol. *Ind Crops Prod* 20:103–110
- Kataoka M, Kotaka A, Thiwhong R, Wada M, Nakamori S, Shimizu S (2004) Cloning and overexpression of the old yellow enzyme gene of *Candida macedoniensis*, and its application to the production of a chiral compound. *J Biotechnol* 114:1–9
- Kiers J, Zeeman AM, Luttkik M, Thiele C, Castrillo JI, Steensma HY, van Dijken JP, Pronk JT (1998) Regulation of alcoholic fermentation in batch and chemostat cultures of *Kluyveromyces lactis* CBS 2359. *Yeast* 14:459–469
- Kim WH, Chung JH, Back JH, Choi J, Cha JH, Koh HY, Han YS (2003) Molecular cloning and characterization of an NADPH quinone oxidoreductase from *Kluyveromyces marxianus*. *J Biochem Mol Biol* 36:442–449
- Kreger-van Rij NJW (1984) The yeasts: a taxonomic study, 3rd edn. Elsevier, Amsterdam
- Künkel W, May R (1976) Alcohol dehydrogenase (ADH) in yeast cells. I. Cytoplasmic, mitochondrial and nuclear ADH in *Saccharomyces carlsbergensis* and *Kluyveromyces fragilis*. *Z Allg Mikrobiol* 16:529–536
- Kurtzman CP (2003) Phylogenetic circumscription of *Saccharomyces*, *Kluyveromyces* and other members of the *Saccharomycetaceae*, and the proposal of the new genera *Lachancea*, *Nakaseomyces*, *Naumovia*, *Vanderwaltozyma* and *Zygorulasporea*. *FEMS Yeast Res* 4:233–245
- Kurtzman CP, Fell JW (eds) (1998) The yeasts: a taxonomic study, 4th edn. Elsevier, Amsterdam
- Kurtzman CP, Robnett CJ (2003) Phylogenetic relationships among yeasts of the ‘*Saccharomyces* complex’ determined from multi-gene sequence analyses. *FEMS Yeast Res* 3:417–432
- Lachance MA (1998) *Kluyveromyces* van der Walt emend. van der Walt. In: Kurtzman CP, Fell JW (eds) The yeasts: a taxonomic study, 4th edn. Elsevier, Amsterdam, pp 227–247
- Lachance MA (2007) Current status of *Kluyveromyces* systematics. *FEMS Yeast Res* 7:642–645
- Ladrière JM, Delcour J, Vandenhoute J (1993) Sequence of a gene coding for a cytoplasmic alcohol dehydrogenase from *Kluyveromyces marxianus* ATCC 12424. *Biochim Biophys Acta* 1173:99–101
- Ladrière JM, Georis I, Guérineau M, Vandenhoute J (2000) *Kluyveromyces marxianus* exhibits an ancestral *Saccharomyces cerevisiae* genome organization downstream of *ADH2*. *Gene* 255:83–91
- Laloux O, Cassart JP, Delcour J, van Beeumen J, Vandenhoute J (1991) Cloning and sequencing of the inulinase gene of *Kluyveromyces marxianus* var. *marxianus* ATCC 12424. *FEBS Lett* 289:64–68
- Leclerc M, Chemardin P, Arnaud A, Ratomahenina R, Galzy P, Gerbaud C, Raynal A, Guérineau M (1987) Comparison of the properties of the purified β -glucosidase from the transformed strain of *Saccharomyces cerevisiae* TYKF2 with that of the donor strain *Kluyveromyces fragilis* Y610. *Biotechnol Appl Biochem* 9:410–422
- Leclercq-Perlat MN, Corrieu G, Spinnler HE (2004) Comparison of volatile compounds produced in model cheese medium deacidified by *Debaryomyces hansenii* or *Kluyveromyces marxianus*. *J Dairy Sci* 87:1545–1550
- Lim J, Yamasaki Y, Suzuki Y, Ozawa J (1980) Multiple forms of endo-polygalacturonase from *Saccharomyces fragilis*. *Agric Biol Chem* 44:473–480
- Llorente B, Malpertuy A, Blandin G, Artiguenave F, Wincker P, Dujon B (2000) Genomic exploration of the hemiascomycetous yeasts: 12. *Kluyveromyces marxianus* var. *marxianus*. *FEBS Lett* 487:71–75
- Lodder J (1970) The yeasts: a taxonomic study, 2nd edn. NHP, Amsterdam
- Lodder J, Kreger-van Rij NJW (1952) The yeasts: a taxonomic study. NHP, Amsterdam
- Love G, Nigam P, Barron N, Singh D, Marchant R, McHale AP (1996) Ethanol production at 45°C using preparations of *Kluyveromyces marxianus* IMB3 immobilized in calcium alginate and kissiris. *Bioproc Eng* 15:275–277
- Love G, Gough S, Brady D, Barron N, Nigam P, Singh D, Marchant R, McHale AP (1998) Continuous ethanol fermentation at 45°C using *Kluyveromyces marxianus* IMB3. Immobilized in calcium alginate and kissiris. *Bioproc Eng* 18:187–189
- Lukondeh T, Ashbolt NJ, Rogers PL (2003a) Evaluation of *Kluyveromyces marxianus* as a source of yeast autolysates. *J Ind Microbiol Biotechnol* 30:52–56
- Lukondeh T, Ashbolt NJ, Rogers PL (2003b) Evaluation of *Kluyveromyces marxianus* FII 510700 grown on a lactose-based medium as a source of a natural bioemulsifier. *J Ind Microbiol Biotechnol* 30:715–720
- Lukondeh T, Ashbolt NJ, Rogers PL (2003c) Confirmation of an alkali-insoluble glucans from *Kluyveromyces marxianus* cultivated

- on a lactose-based medium. *World J Microbiol Biotechnol* 19: 349–355
- Mahoney RR, Nickerson TA, Whitaker JR (1975) Selection of strain, growth conditions and extraction procedures for optimum production of lactase from *Kluyveromyces fragilis*. *J Dairy Sci* 58:1620–1629
- Majumdar S, Bhattacharjee H, Bhattacharyya D, Bhaduri A (1998) UDP-galactose 4-epimerase from *Kluyveromyces fragilis*: reconstitution of holoenzyme structure after dissociation with parachloromercuribenzoate. *Eur J Biochem* 257:427–433
- Margaritis A, Bajpai P (1982) Direct fermentation of D-xylose to ethanol by *Kluyveromyces marxianus* strains. *Appl Environ Microbiol* 44:1039–1041
- Margaritis A, Bajpai P (1983) Effect of sugar concentration in Jerusalem artichoke extract on *Kluyveromyces marxianus* growth and ethanol production. *Appl Environ Microbiol* 45:723–725
- Martins DB, de Souza CG Jr, Simões DA, de Morais MA Jr (2002) The b-galactosidase activity in *Kluyveromyces marxianus* CBS6556 decreases by high concentrations of galactose. *Curr Microbiol* 44:379–382
- Marwaha SS, Kennedy JF, Sehgal VK (1988) Simulation of process conditions of continuous ethanol fermentation of whey permeate using alginate entrapped *Kluyveromyces marxianus* NCYC-179 cells in a packed-bed reactor system. *Proc Biochem* 23:17–22
- Medeiros ABP, Pandey A, Freitas RJS, Christen P, Soccol CR (2000) Optimization of the production of aroma compounds by *Kluyveromyces marxianus* in solid-state fermentation using factorial design and response surface methodology. *Biochem Eng J* 6:33–39
- Medeiros ABP, Pandey A, Christen P, Fontoura PSG, de Freitas RJS, Soccol CR (2001) Aroma compounds produced by *Kluyveromyces marxianus* in solid state fermentation on a packed bed column bioreactor. *World J Microbiol Biotechnol* 17:767–771
- Meilhoc E, Masson JM, Teissie J (1990) High efficiency transformation of intact yeast cells by electric field pulses. *J Biotechnol* 8:223–227
- Molnár O, Prillinger H, Lopandic K, Weigang F, Staudacher E (1996) Analysis of coenzyme Q systems, monosaccharide patterns of purified cell walls, and RAPD-PCR patterns in the genus *Kluyveromyces*. *Antonie van Leeuwenhoek* 70:67–78
- Mukherji S, Bhaduri A (1992) An essential histidine residue for the activity of UDPglucose 4-epimerase from *Kluyveromyces fragilis*. *J Biol Chem* 267:11709–11713
- Neves L, Oliveira R, Lucas C (2004) Yeast orthologues associated with glycerol transport and metabolism. *FEMS Yeast Res* 5:51–62
- Nguyen TH, Fleet GH, Rogers PL (1998) Composition of the cell walls of several yeast species. *Appl Microbiol Biotechnol* 50:206–212
- Nigam P, Banat IM, Singh D, McHale AP, Marchant R (1997) Continuous ethanol production by thermotolerant *Kluyveromyces marxianus* IMB3 yeast immobilized on mineral kisseris at 45°C. *World J Microbiol Biotechnol* 13:283–288
- Nilsson U, Barron N, McHale L, McHale AP (1995) The effects of phosphoric and pretreatment on conversion of cellulose to ethanol at 45°C using the thermotolerant yeast *Kluyveromyces marxianus* IMB3. *Biotechnol Lett* 17:985–988
- Nolan AM, Barron N, Brady D, McAree T, Smith D, McHale L, McHale AP (1994) Ethanol production at 45°C by an alginate-immobilized strain of *Kluyveromyces marxianus* following growth on glucose-containing media. *Biotechnol Lett* 16:849–852
- Nunes MFA, Massaguer S, Alegre RM (1993) Produção e propriedades de b-galactosidase de *Kluyveromyces marxianus* NRRL Y-2415. *Rev Farm Bioquim Univ S Paulo* 29:25–30
- Oberyé EH, Maurer K, Mager WH, Planta RJ (1993) Structure of the ABF1-homologue from *Kluyveromyces marxianus*. *Biochim Biophys Acta* 1173:233–236
- O'Shea DG, Walsh PK (2000) The effect of culture conditions on the morphology of the dimorphic yeast *Kluyveromyces marxianus* var. *marxianus* NRRLy 2415: a study incorporating image analysis. *Appl Microbiol Biotechnol* 53:316–322
- Panuwatsuk W, da Silva NA (2002) Evaluation of pKD1-based plasmid systems for heterologous protein production in *Kluyveromyces lactis*. *Appl Microbiol Biotechnol* 58:195–201
- Parekh S, Margaritis A (1985) Inulinase (b-fructofuranosidase) production by *Kluyveromyces marxianus* in batch culture. *Appl Microbiol Biotechnol* 22:446–448
- Pecota DC, da Silva NA (2005) Evaluation of the tetracycline promoter system for regulated gene expression in *Kluyveromyces marxianus*. *Biotechnol Bioeng* 92:117–123
- Pecota DC, Rajgarhia V, da Silva NA (2007) Sequential gene integration for the engineering of *Kluyveromyces marxianus*. *J Biotechnol* 127:408–416
- Penman CS, Duffus JH (1974) Ergosterol is the only sterol in *Kluyveromyces fragilis*. *Antonie van Leeuwenhoek* 40:529–531
- Pessoa Jr A, Vitolo M (1999) Inulinase from *Kluyveromyces marxianus*: culture medium composition and enzyme extraction. *Braz J Chem Eng* 16:237–245
- Pinheiro R, Belo I, Mota M (2000) Air pressure effects on biomass yield of two different *Kluyveromyces* strains. *Enzyme Microb Technol* 26:756–762
- Pinheiro R, Belo I, Mota M (2002) Oxidative stress response of *Kluyveromyces marxianus* to hydrogen peroxide, paraquat and pressure. *Appl Microbiol Biotechnol* 58:842–847
- Porro D, Sauer M, Branduardi P, Mattanovich D (2005) Recombinant protein production in yeasts. *Mol Biotechnol* 31:245–259
- Postma E, van der Broek PJA (1990) Continuous-culture study of the regulation of glucose and fructose transport in *Kluyveromyces marxianus* CBS 6556. *J Bacteriology* 172:2871–2876
- Prudêncio C, Sansonetty F, Sousa MJ, Côrte-Real M, Leão C (2000) Rapid detection of efflux pumps and their relation with drug resistance in yeast cells. *Cytometry* 39:26–35
- Queirós O, Casal M, Althoff S, Moradas-Ferreira P, Leão C (1998) Isolation and characterization of *Kluyveromyces marxianus* mutants deficient in malate transport. *Yeast* 14:401–407
- Rajoka MI, Khan S, Shahid R (2003) Kinetics and regulation studies of the production of b-galactosidase from *Kluyveromyces marxianus* grown on different substrates. *Food Technol Biotechnol* 41:315–320
- Rajoka MI, Latif F, Khan S, Shahid R (2004) Kinetics of improved productivity of b-galactosidase by a cycloheximide-resistant mutant of *Kluyveromyces marxianus*. *Biotechnol Lett* 26:741–746
- Ramirez-Zavala B, Mercado-Flores Y, Hernandez-Rodriguez C, Villatanaca L (2004a) Purification and characterization of a lysine aminopeptidase from *Kluyveromyces marxianus*. *FEMS Microbiol Lett* 235:369–375
- Ramirez-Zavala B, Mercado-Flores Y, Hernandez-Rodriguez C, Villatanaca L (2004b) Purification and characterization of a serine carboxypeptidase from *Kluyveromyces marxianus*. *Int J Food Microbiol* 91:245–252
- Ray S, Mukherji S, Bhaduri A (1995) Two tryptophans at the active site of UDP-glucose 4-epimerase from *Kluyveromyces fragilis*. *J Biol Chem* 270:11383–11390
- Raynal A, Guerineau M (1984) Cloning and expression of the structural gene for b-glucosidase of *Kluyveromyces fragilis* in *Escherichia coli* and *Saccharomyces cerevisiae*. *Mol Gen Genet* 195:108–115
- Raynal A, Gerbaud C, Francingues MC, Guerineau M (1987) Sequence and transcription of the b-glucosidase gene of *Kluyveromyces fragilis* cloned in *Saccharomyces cerevisiae*. *Curr Genet* 12:175–184
- Rech R, Cassini CF, Secchi AR, Ayub MAZ (1999) Utilization of protein-hydrolyzed chesse whey for the production of b-galactosidase by *Kluyveromyces marxianus*. *J Ind Microbiol Biotechnol* 23:91–96

- Ribeiro O, Gombert AK, Teixeira JA, Domingues L (2007) Application of the Cre-loxP system for multiple gene disruption in the yeast *Kluyveromyces marxianus*. *J Biotechnol* 131:20–26
- Riordan C, Love G, Barron N, Nigam P, Marchant R, McHale L, McHale AP (1996) Production of ethanol from sucrose at 45°C by alginate-immobilized preparations of the thermotolerant yeast strain *Kluyveromyces marxianus* IMB3. *Biores Technol* 55:171–173
- Romanos MA, Scorer CA, Clare JJ (1992) Foreign gene expression in yeast: a review. *Yeast* 8:423–488
- Rosa FM, Sa-Correia I (1992) Ethanol tolerance and activity of plasma membrane ATPase in *Kluyveromyces marxianus* and *Saccharomyces cerevisiae*. *Enzyme Microb Technol* 14:23–27
- Rouwenhorst RJ, Visser LE, van der Baan AA, Scheffers WA, van Dijken JP (1988) Production, distribution, and kinetic properties of inulinase in continuous culture of *Kluyveromyces marxianus* CBS 6556. *Appl Environ Microbiol* 54:1131–1137
- Rouwenhorst RJ, Hensing M, Verbakel J, Scheffers WA, van Dijken JP (1990a) Structure and properties of the extracellular inulinase of *Kluyveromyces marxianus* CBS 6556. *Appl Environ Microbiol* 56:3337–3345
- Rouwenhorst RJ, Ritmeester WS, Scheffers WA, van Dijken JP (1990b) Localization of inulinase and invertase in *Kluyveromyces* species. *Appl Environ Microbiol* 56:3329–3336
- Sakanaka K, Yan W, Kishida M, Sakai T (1996) Breeding a fermentative yeast at high temperature using protoplast fusion. *J Ferment Bioeng* 81:104–108
- Schaffrath R, Breunig KD (2000) Genetics and molecular physiology of the yeast *Kluyveromyces lactis*. *Fung Genet Biol* 30:173–190
- Scharpf LG, Seitz EW, Morris JA, Farbood MI (1986) Generation of flavor and odor compounds through fermentation processes. In: Parliament TH, Croteau R (eds) *Biogenesis of aroma*, American Chemical Society, Washington, DC, vol. 317, pp 323–346
- Schultz N, Chang L, Hauck A, Reuss M, Syldatk C (2006) Microbial production of single-cell protein from deproteinized whey concentrates. *Appl Microbiol Biotechnol* 69:515–520
- Schwan RF, Rose AH (1994) Polygalacturonase production by *Kluyveromyces marxianus*: effect of medium composition. *J Appl Bacteriol* 76:62–67
- Schwan RF, Cooper RM, Wheals AE (1997) Endopolygalacturonase secretion by *Kluyveromyces marxianus* and other cocoa pulp-degrading yeasts. *Enzyme Microb Technol* 21:234–244
- Serrat M, Bermudez RC, Villa TG (2004) Polygalacturonase and ethanol production in *Kluyveromyces marxianus*—potential use of polygalacturonase in foodstuffs. *Appl Biochem Biotechnol* 117:49–64
- Šiekštelė R, Bartkevičiūtė D, Sasnauskas K (1999) Cloning, targeted disruption and heterologous expression of the *Kluyveromyces marxianus* endopolygalacturonase gene (*EPGI*). *Yeast* 15:311–322
- Skountzou P, Soupioni M, Bekatorou A, Kanellaki M, Koutinas AA, Marchant R, Banat IM (2003) Lead(II) uptake during baker's yeast production by aerobic fermentation of molasses. *Proc Biochem* 38:1479–1482
- Sonawat HM, Agrawal A, Dutta SM (1981) Production of β-galactosidase from *Kluyveromyces fragilis* grown on whey. *Folia Microbiol* 26:370–376
- Souciet J-L, Artiguenave MAF, Blandin G, Bolotin-Fukuhara M, Bon E, Brottier P et al (2000) Genomic exploration of the hemiascomycetous yeasts: 1. A set of yeast species for molecular evolution studies. *FEBS Letters* 487:3–12
- Stambuk BU, Franden MA, Singh A, Zhang M (2003) D-Xylose transport by *Candida succiphila* and *Kluyveromyces marxianus*. *Appl Biochem Biotechnol* 108:255–263
- Steenma HY, Ter Linde JJ (2001) Plasmids with the Cre-recombinase and the dominant nat marker, suitable for use in prototrophic strains of *Saccharomyces cerevisiae* and *Kluyveromyces lactis*. *Yeast* 18:469–472
- Steenma HY, de Jongh FCM, Linnekamp M (1988) The use of electrophoretic karyotypes in the classification of yeasts: *Kluyveromyces marxianus* and *K. lactis*. *Curr Genet* 14:311–317
- Stephanopoulos GN, Aristidou AA, Nielsen J (1998) *Metabolic Engineering Principles and Methodologies*. Academic Press, San Diego
- Sukroongreung S, Schappert KT, Khachatourians GG (1984) Survey of sensitivity of twelve yeast genera toward T-2 toxin. *Appl Environ Microbiol* 48:416–419
- Tin CSF, Mawson AJ (1993) Ethanol production from whey in a membrane recycle bioreactor. *Proc Biochem* 28:217–221
- Ternan NG, McMullan G (2000) The utilization of 4-aminobutylphosphonate as sole nitrogen source by a strain of *Kluyveromyces fragilis*. *FEMS Microbiol Lett* 184:237–240
- Ternan NG, McMullan G (2002) Iminodiacetate and nitrilotriacetate degradation by *Kluyveromyces marxianus* IMB3. *Biochem Biophys Res Comm* 290:802–805
- Toyoda Y, Sy J (1984) Purification and phosphorylation of fructose-1,6-bisphosphatase from *Kluyveromyces fragilis*. *J Biol Chem* 259:8718–8723
- van den Berg JA, van der Laken KJ, van Ooyen AJJ, Renniers TCHM, Rietveld K et al (1990) *Kluyveromyces* as a host for heterologous gene expression. Expression and secretion of prochymosin. *Bio/Techno* 8:135–139
- van den Broek PJ, de Bruijne AW, van Steveninck J (1987) The role of ATP in the control of H⁺-galactoside symport in the yeast *Kluyveromyces marxianus*. *Biochem J* 242:729–734
- van der Walt JP (1956) *Kluyveromyces*—a new yeast genus of the Endomycetales. *Antonie van Leeuwenhoek* 22:265–272
- van der Walt JP (1970) *Kluyveromyces* van der Walt emend. van der Walt. In: Lodder J (ed) *The yeasts: a taxonomic study*, 2nd edn. NHPG, Amsterdam, pp 316–378
- van der Walt JP, Johannsen E (1984) *Kluyveromyces* van der Walt emend. van der Walt. In: Kreger-van Rij NJW (ed) *The yeasts: a taxonomic study*, 3rd edn. Elsevier, Amsterdam, pp 224–251
- van Dijken JP, Weusthuis RA, Pronk JT (1993) Kinetics of growth and sugar consumption in yeasts. *Antonie van Leeuwenhoek* 63:343–352
- van Dijken JP, Bauer J, Brambilla L et al (2000) An interlaboratory comparison of physiological and genetic properties of four *Saccharomyces cerevisiae* strains. *Enz Microb Technol* 26:706–714
- van Leeuwen CC, Postma E, van den Broek PJ, van Steveninck J (1991) Proton-motive force-driven D-galactose transport in plasma membrane vesicles from the yeast *Kluyveromyces marxianus*. *J Biol Chem* 266:12146–12151
- van Ooyen AJ, Dekker P, Huang M, Olsthoorn MM, Jacobs DI, Colussi PA, Taron CH (2006) Heterologous protein production in the yeast *Kluyveromyces lactis*. *FEMS Yeast Res* 6:381–392
- van Urk H, Voll WSL, Scheffers WA, van Dijken JP (1990) Transient-state analyses of metabolic fluxes in Crabtree-positive and Crabtree-negative yeasts. *Appl Environ Microbiol* 56:281–287
- Verduyn C, Postma E, Scheffers WA, van Dijken JP (1992) Effect of benzoic acid on metabolic fluxes in yeasts: a continuous-culture study on the regulation of respiration and alcoholic fermentation. *Yeast* 8:501–517
- Verzotti E, Geymonat M, Valetti F, Lanzetti L, Giunta C (1994) In the budding yeast *Kluyveromyces marxianus*, adenylate cyclase is regulated by Ras protein(s) in vitro. *Yeast* 10:993–1001
- Viegas CA, Rosa MF, Sá-Correia I, Novais JM (1989) Inhibition of yeast growth by octanoic and decanoic acids produced during ethanolic fermentation. *Appl Environ Microbiol* 55:21–28
- Visser W, Scheffers WA, Batenburg-van der Vegte WH, van Dijken JP (1990) Oxygen requirements of yeasts. *Appl Environ Microbiol* 56:3785–3792

- Yoda K, Ko JH, Nagamatsu T, Lin Y, Kaibara C, Kawada T, Tomishige N, Hashimoto H, Noda Y, Yamasaki M (2000) Molecular characterization of a novel yeast cell-wall acid phosphatase cloned from *Kluyveromyces marxianus*. *Biosci Biotechnol Biochem* 64:142–148
- Ward C, Nolan AM, O'Hanlon F, McAree T, Barron N, McHale L, McHale AP (1995) Production of ethanol at 45°C on starch-containing media by mixed cultures of the thermotolerant, ethanol-producing yeast *Kluyveromyces marxianus* IMB3 and the thermophilic filamentous fungus *Talaromyces emersonii* CBS 813.70. *Appl Microbiol Biotechnol* 43:408–411
- Wésolowski-Louvel M, Breunig KD, Fukuhara H (1996) *Kluyveromyces lactis*: genetics, biochemistry and molecular biology of non-conventional yeast. Springer-Verlag, Berlin Heidelberg New York
- Welsh FW, Murray WD, Williams RE (1989) Microbiological and enzymatic production of flavor and fragrance chemicals. *Crit Rev Biotechnol* 9:105–169
- Wimborne MP, Rickard PAD (1978) Pectinolytic activity of *Saccharomyces fragilis* cultured in controlled environments. *Biotechnol Bioeng* 20:231–242
- Wittmann C, Hans M, Bluemke W (2002) Metabolic physiology of aroma-producing *Kluyveromyces marxianus*. *Yeast* 19:1351–1363
- Wolf K, Breunig K, Barth G (eds) (2003) Non-conventional yeasts in genetics, biochemistry and biotechnology: practical protocols. Springer-Verlag, Berlin Heidelberg New York
- Workman WE, Day DF (1984) The cell wall-associated inulinase of *Kluyveromyces fragilis*. *Antonie van Leeuwenhoek* 50: 349–353
- Yoshida Y, Yokoi W, Wada Y, Ohishi K, Ito M, Sawada H (2004) Potent hypocholesterolemic activity of the yeast *Kluyveromyces marxianus* YIT 8292 in rats fed a high cholesterol diet. *Biosci Biotechnol Biochem* 68:1185–1192
- Yoshida Y, Yokoi W, Ohishi K, Ito M, Naito E, Sawada H (2005) Effects of the cell wall of *Kluyveromyces marxianus* YIT 8292 on the plasma cholesterol and fecal sterol excretion in rats fed on a high-cholesterol diet. *Biosci Biotechnol Biochem* 69:714–723
- Zhang J, Yuan H, Wen T, Xu F, Di Y, Huo K, Li YY (2003) Cloning of the *KcURA3* gene and development of a transformation system for *Kluyveromyces cicerisporus*. *Appl Microbiol Biotechnol* 62: 387–391